

## Date:     /     /     Page:     /

PTL diagnosis - Date: \_\_\_\_/\_\_\_\_/\_\_\_\_ Time: \_\_\_\_:\_\_\_\_

**Gestational age:** \_\_\_\_ wks + \_\_\_\_ days

ALERT ← FIRST STAGE → SECOND STAGE →

[illegible]

INSTRUCTIONS: CIRCLE ANY OBSERVATION MEETING THE CRITERIA IN THE 'ALERT' COLUMN, ALERT THE SENIOR MIDWIFE OR DOCTOR AND RECORD THE ASSESSMENT AND ACTION TAKEN. IF LABOUR EXTENDS BEYOND 12H, PLEASE CONTINUE ON A NEW LABOUR CARE GUIDE FOR PTL/PTB. Abbreviations: Y – Yes, N – No, D – Declined, U – Unknown, E – Early, L – Late, V – Variable, I – Intact, C – Clear, M – Meconium, B – Blood, P – Purulent, OA – Occiput Anterior, OP – Occiput Posterior, OT – Occiput Transverse, B – Breech, TL – Transverse lie, P+ – Protein, A+ – Acetone

Developed for the WHO Implementation Research to Scale-up and Evaluate the Impact of Antenatal Corticosteroids on Preterm Newborn Outcomes (ACS-IR) and should only be used in the context of research. The Labour Care Guide for PTL should be used in conjunction with the Quick Guide. Responsibility for the interpretation and use of the material lies with the reader. In no event shall the WHO be liable for damages arising from its use.